

KF/Al₂O₃: As a Solid Phase and Recyclable Basic Catalyst for Synthesis Mono and Bis Pyrimidine Derivatives

Abaszadeh, Mehdi *⁺

Pharmaceutics Research Center, Institute of Neuropharmacology, Kerman University of Medical Sciences, Kerman, I.R. IRAN

Seifi, Mohammad

Department of Chemistry, Faculty of Sciences, Najafabad Branch, Islamic Azad University, Najafabad, I.R. IRAN

ABSTRACT: *KF/Al₂O₃ as a green and efficient catalyst has been used for the synthesis of mono and bis pyrimidine derivatives via three-component reaction of amidines, malononitrile, and aldehydes in EtOH at reflux. The great advantage of this catalyst is the ease of handling. KF/Al₂O₃ can be used and removed by filtration, avoid cumbersome aqueous workups and decrease solvent waste handling issues. The additional benefit is achieved by taking advantage of the strongly basic nature of KF/Al₂O₃, which allows replacing organic bases in the reaction. A number of advantages are with this method such as no special handling necessity of catalyst, easy monitoring of reaction process, convenient workup procedure and high yields in short reaction times.*

KEYWORDS: *KF/Al₂O₃; Solid phase and recyclable basic catalyst; Mono and bis pyrimidine derivatives; Three-component reaction.*

INTRODUCTION

Recently, solid phase reagents have been preferred for carrying out various chemical transformations due to the good activation of adsorbed compounds, increase in reaction rate, selectivity, milder reaction conditions and easier workup [1]. Potassium fluoride on alumina is one of the most interesting of these reagents because it has surface properties which suggest that very rich organic reactions may occur there. KF/Al₂O₃ is an inexpensive and commercially available reagent which due to its strongly basic nature it has been used as a replacement for organic bases in a number of organic reactions, such as epoxidation [2], Michael addition [3], aldol condensation [4], rearrangement processes [5], cycloaddition reactions [6, 7], multicomponent reactions

[8-11], for conversion of aldoximes into nitriles [12], α -phenylselenenylation of aldehydes and ketones [13] and synthesis of nitriles from aldehydes [14].

Multi-Component Reactions (MCRs) are efficient and effective methods to synthesis complex molecules with special synthetic yield, frequently with high stereoselectivity, easily accessible reactants [15-17]. The characteristic aspect of MCRs is that the final products contain almost all portions of substrates, generating almost no by-products, which make MCRs an extremely ideal and eco-friendly reaction system to produce many organic compounds [18-21].


It is not admiring that the synthesis of polyfunctionalized heterocyclic compounds has received substantial attention

* To whom correspondence should be addressed.

+ E-mail: abaszadeh@kmu.ac.ir

1021-9986/2017/1/35

9/\$/5.90



Scheme 1: Pyrimidine in nature and in synthetic compounds.

because polyfunctionalized heterocyclic compounds play important roles in the synthesis of a large number of biologically active molecules [22-25]. In recent years, a great deal of research in polyfunctionalized heterocyclic compounds paid to the synthesis of bis-heterocyclic compounds which show various biological activities, such as antibacterial, fungicidal, tuberculostatic, antiamebic properties [26-28]. Among those, pyrimidine derivatives have recovered much attention in recent years due to their notable molecular structures and their important biological activities.

Pyrimidine is one of the most common *N*-heteroaromatic compounds containing two nitrogen atoms has widely appeared in nature as substituted and ring fused compounds and derivatives, including the thiamine (vitaminB1) and alloxan (Scheme 1) [29]. Several pyrimidines mainly uracil, thymine, and cytosine (Scheme 1) have been isolated from the nucleic acid hydrolyses [30]. It is also found in many synthetic compounds such as veranal which is used as hypnotics [31] and the HIV drug, zidovudine [32] (Scheme 1). In addition to this, pyrimidines ring demonstrate a diverse array of biological and pharmacological activities including, antibacterial [33], analgesics [34], antipyretics [35], anti-inflammatory [36], antifungal [37], and anticancer [38] properties.

There are various catalytic system to synthesis of pyrimidine derivatives by MCRs, such as sodium

acetate [39, 40], potassium carbonate in the presence of tetrabutylammonium bromide (TBAB) [41], high surface area MgO (HSA-MgO) [42], copper oxide nanocatalyst [43] and $\text{Bi}(\text{NO}_3)_3 \cdot 5\text{H}_2\text{O}$ and a catalytic amount of Et_3N [44]. But these systems have some disadvantage like the use of toxic organic solvents, costly catalysts, the existence of transition metals, time-consuming reaction, and low yield. Herein we used $\text{KF}/\text{Al}_2\text{O}_3$ as catalyst for the synthesis of 4-amino-5-pyrimidine-carbonitrile derivatives to remove the above-mentioned drawbacks. We consider the synthesis of compound 2,4-diamino-6-phenylpyrimidine-5-carbonitrile (**4a**) for a representative example to show the advantage of this work in comparison with previously reported procedures. As shown in Table 1, our catalyst produces compound **4a** with high yield in a short time than others. Also, $\text{KF}/\text{Al}_2\text{O}_3$ is a recyclable catalyst and ethanol was used as a green solvent. Moreover, $\text{KF}/\text{Al}_2\text{O}_3$ is more stable in air and non-toxic compared with other catalysts.

EXPERIMENTAL SECTION

Melting points were measured on a Electrothermal-9100 apparatus and are uncorrected. IR spectra were recorded on a Bruker FT-IR Tensor 27 infrared spectrophotometer. ^1H NMR and spectra were recorded on an Avance III 400 MHz Bruker spectrometer. ^{13}C NMR spectra were recorded on the same instruments at 100 MHz using TMS as an internal standard respectively.

Table 1: Comparison results of KF/Al₂O₃ with other catalysts reported in the literature for the synthesis of compound 4a.

Entry	Catalyst	Condition	Recyclability	Time (min)	Yield (%)
1	CH ₃ COONa	H ₂ O/C ₂ H ₅ OH, reflux	No	300	81 [39]
2	K ₂ CO ₃ /TBAB	H ₂ O, reflux	No	240	63 [41]
3	HSA-MgO	CH ₃ CN, reflux	No	15	88 [42]
4	Nano-CuO	H ₂ O, r.t.	Yes (5 cycles)	15	93 [43]
5	Bi(NO ₃) ₃ .5H ₂ O/Et ₃ N	CH ₃ CN, reflux	No	4	83 [44]
6	KF/Al ₂ O ₃	EtOH, reflux	Yes (4 cycles)	5	94 [This work]

Elemental analyses were performed using a Heracus CHN-O-Rapid analyzer.

Preparation of KF/Al₂O₃

A mixture of KF (5 mmol) and Al₂O₃ (5 mmol) was ground vigorously in a mortar to give the KF/Al₂O₃ reagent as a white powder (0.800 g) [45].

General procedure for the preparation of mono and bis 4-amino-5-pyrimidine-carbonitrile derivatives (4a-v)

A mixture of amidines (**1a-c**) (2 mmol), malononitrile (**2**) (2 mmol), aromatic aldehydes (**3a-g**) (2 mmol) (if terephthalaldehyde or isophthalaldehyde (**3h, 3i**) (1 mmol)), and KF/Al₂O₃ (30 mol%) in ethanol (10 mL) was refluxed for the time reported in Table 3 (the progress of the reaction being monitored by TLC and was used hexane/ethyl acetate as an eluent). After completion of the reaction, the catalyst was separated by a simple filtration from the reaction mixture. Then the reaction mixture was poured into ice-cold water; the crude product was filtered, dried, and recrystallized from ethanol.

2,4-Diamino-6-(furan-2-yl)pyrimidine-5-carbonitrile (4k)

Brown powders; IR (KBr, ν max/cm⁻¹): 3392-3200 (2NH₂), 2192 (CN), 1619 (C=N), 1587 (C=C). ¹H NMR (400 MHz, DMSO-d₆): 8.11 (d, 1H, ³J_{HH}=8 Hz, CH-Ar),

8.01-7.59 (m, 4H, CH-Ar, NH₂), 7.37 (s, 2H, NH₂). ¹³C NMR (100 MHz, DMSO-d₆): 170.14, 166.08, 163.58, 151.22, 142.35, 131.62, 126.14, 114.31 (CN), 84.69 (C5). Anal. calcd. for C₉H₇N₅O: C, 53.73; H, 3.51; N, 34.81 %. Found: C, 53.74; H, 3.33; N, 34.63 %.

4-Amino-6-(furan-2-yl)-2-phenylpyrimidine-5-carbonitrile (4l)

Orange powders; IR (KBr, ν max/cm⁻¹): 3392, 3216 (NH₂), 2176 (CN), 1607 (C=N), 1593 (C=C). ¹H NMR (400 MHz, DMSO-d₆): 8.39 (d, 2H, ³J_{HH}=8 Hz, CH-Ar), 8.07 (d, 1H, ³J_{HH}=8 Hz, CH-Ar), 7.78-6.87 (m, 7H, CH-Ar, NH₂). ¹³C NMR (100 MHz, DMSO-d₆): 170.69, 166.34, 164.27, 152.13, 141.47, 135.65, 132.27, 128.36, 127.25, 122.58, 121.44, 114.15 (CN), 85.47 (C5). Anal. calcd. for C₁₅H₁₀N₄O: C, 68.69; H, 3.84; N, 21.36 %. Found: C, 68.51; H, 3.66; N, 21.17 %.

4-Amino-6-(furan-2-yl)-2-methylpyrimidine-5-carbonitrile (4m)

Red powders; IR (KBr, ν max/cm⁻¹): 3360, 3312 (NH₂), 2176 (CN), 1619 (C=N), 1584 (C=C). ¹H NMR (400 MHz, DMSO-d₆): 8.24 (d, 1H, ³J_{HH}=8 Hz, CH-Ar), 8.09-7.89 (m, 2H, CH-Ar), 7.59 (s, 2H, NH₂), 2.47 (s, 3H, CH₃). ¹³C NMR (100 MHz, DMSO-d₆): 170.02, 167.65, 163.58, 151.14, 136.20, 135.17, 128.11, 114.35 (CN), 84.78 (C5), 24.93 (CH₃). Anal. calcd. for

$C_{10}H_8N_4O$: C, 59.99; H, 4.03; N, 27.99 %. Found: C, 59.81; H, 3.85; N, 27.80 %.

2,4-Diamino-6-(thiophen-2-yl)pyrimidine-5-carbonitrile (4n)

Red powders; IR (KBr, ν max/cm⁻¹): 3328-3200 (2NH₂), 2192 (CN), 1616 (C=N), 1590 (C=C). ¹H NMR (400 MHz, DMSO-d₆): 7.97-7.39 (m, 5H, CH-Ar, NH₂), 7.01 (s, 2H, NH₂). ¹³C NMR (100 MHz, DMSO-d₆): 170.24, 165.88, 163.47, 141.40, 132.27, 129.25, 126.01, 113.89 (CN), 84.57 (C5). Anal. calcd. for C₉H₇N₃S: C, 49.76; H, 3.25; N, 32.24 %. Found: C, 49.56; H, 3.06; N, 32.25 %.

4-Amino-2-methyl-6-(thiophen-2-yl)pyrimidine-5-carbonitrile (4p)

Orange powders; IR (KBr, ν max/cm⁻¹): 3344, 3232 (NH₂), 2176 (CN), 1607 (C=N), 1593 (C=C). ¹H NMR (400 MHz, DMSO-d₆): 8.01 (d, 1H, ³J_{HH}=8 Hz, CH-Ar), 7.92 (d, 1H, ³J_{HH}=8 Hz, CH-Ar), 7.68-7.35 (m, 3H, CH-Ar, NH₂), 2.27 (s, 3H, CH₃). ¹³C NMR (100 MHz, DMSO-d₆): 171.25, 166.98, 163.22, 144.75, 135.22, 129.67, 121.33, 114.25 (CN), 85.34 (C5), 25.12 (CH₃). Anal. calcd. for C₁₀H₈N₄S: C, 55.54; H, 3.73; N, 25.91 %. Found: C, 55.36; H, 3.55; N, 25.73 %.

6,6'-(1,4-Phenylene) bis(2,4-diaminopyrimidine-5-carbonitrile) (4q)

Orange powders; IR (KBr, ν max/cm⁻¹): 3344-3200 (2NH₂), 2192 (CN), 1622 (C=N), 1580 (C=C). ¹H NMR (400 MHz, DMSO-d₆): 7.92-7.06 (m, 8H, 2NH₂, CH-Ar), 6.53 (s, 4H, 2NH₂). ¹³C NMR (100 MHz, DMSO-d₆): 168.19, 164.77, 159.36, 137.28, 122.29, 115.07 (CN), 84.08 (C5). Anal. calcd. For C₁₆H₁₂N₁₀: C, 55.81; H, 3.51; N, 40.68 %. Found: C, 55.62; H, 3.33; N, 40.50 %.

6,6'-(1,3-Phenylene) bis(2,4-diaminopyrimidine-5-carbonitrile) (4t)

Yellow powders; IR (KBr, ν max/cm⁻¹): 3344-3200 (2NH₂), 2192 (CN), 1616 (C=N), 1558 (C=C). ¹H NMR (400 MHz, DMSO-d₆): 7.89-7.23 (m, 8H, 2NH₂, CH-Ar), 6.67 (s, 4H, 2NH₂). ¹³C NMR (100 MHz, DMSO-d₆): 168.10, 164.70, 159.94, 137.19, 126.63, 114.87 (CN), 84.13 (C5). Anal. calcd. For C₁₆H₁₂N₁₀: C, 55.81; H, 3.51; N, 40.68 %. Found: C, 55.63; H, 3.32; N, 40.50 %.

6,6'-(1,3-Phenylene) bis(4-amino-2-phenylpyrimidine-5-carbonitrile) (4u)

Yellow powders; IR (KBr, ν max/cm⁻¹): 3360, 3216 (NH₂), 2176 (CN), 1619 (C=N), 1574 (C=C). ¹H NMR (400 MHz, DMSO-d₆): 8.45 (s, 4H, 2NH₂), 7.79-7.12 (m, 14H, CH-Ar). ¹³C NMR (100 MHz, DMSO-d₆): 168.69, 166.02, 162.35, 141.80, 136.58, 133.25, 129.07, 128.67, 126.38, 125.94, 121.25, 115.01 (CN), 84.20 (C5). Anal. calcd. For C₂₈H₁₈N₈: C, 72.09; H, 3.89; N, 24.02 %. Found: C, 71.90; H, 3.71; N, 23.83 %.

6,6'-(1,3-Phenylene) bis(4-amino-2-methylpyrimidine-5-carbonitrile) (4v)

Yellow powders; IR (KBr, ν max/cm⁻¹): 3344, 3200 (NH₂), 2176 (CN), 1628 (C=N), 1545 (C=C). ¹H NMR (400 MHz, DMSO-d₆): 8.84 (s, 4H, 2NH₂), 8.04-7.75 (m, 4H, CH-Ar), 2.47 (s, 3H, CH₃). ¹³C NMR (100 MHz, DMSO-d₆): 168.10, 164.77, 161.01, 139.97, 137.38, 128.54, 126.63, 114.49 (CN), 84.25 (C5), 26.49 (CH₃). Anal. calcd. For C₁₈H₁₄N₈: C, 63.15; H, 4.12; N, 32.73 %. Found: C, 62.96; H, 3.94; N, 32.54 %.

RESULTS AND DISCUSSION


In continuation of our previous works on environmentally friendly multi-component reactions [46-49], we have synthesized a series of mono and bis 4-amino-5-pyrimidine-carbonitrile derivatives (**4a-v**) via three-component reaction of amidines (**1a-c**), malononitrile (**2**) and aldehydes (**3a-i**) in presence of KF/Al₂O₃ in EtOH at reflux (Scheme 2).

In order to optimize the reaction conditions, we used some polar and nonpolar solvents in the three-component reaction of guanidine **1a**, malononitrile **2** and benzaldehyde **3a** in the presence of 30 mol% KF/Al₂O₃ as model reactions to examine the effects of solvent for preparing compound **4a**. In each case, the substrates were mixed together with 30 mol% KF/Al₂O₃ agitated with 10 mL solvent under reflux. The results are shown in Table 2 (entry 1-4). The best results in terms of reaction time and yield of the desired product **4a** was obtained when the reaction was carried out in ethanol.

We also optimized the quantity of catalysts. The best results were obtained when the reactions were carried out in the presence of 30 mol% KF/Al₂O₃. The results are shown in Table 2 (entry 5-8).

Table 2: Optimization of the model reaction between guanidine 1a, malononitrile 2 and benzaldehyde 3a.

Entry	Solvent	Catalyst (mol %)	Time (min)	Yield (%)
1	EtOH*	30	5	94
2	CH ₃ CN	30	30	85
3	EtOAc	30	45	82
4	Toluene	30	60	55
5	EtOH	10	15	88
6	EtOH	20	10	90
7	EtOH	25	7	92
8	EtOH	35	5	94
9 ^a	EtOH	30	7	92
10 ^b	EtOH	30	10	91
11 ^c	EtOH	30	15	90
12 ^d	EtOH	30	15	88

**Scheme 2: Synthesis of mono and bis 4-amino-5-pyrimidine-carbonitrile derivatives.**

We also attempted to reuse the catalyst. After each cycle, the catalyst was recovered by simple filtration, washed with hot ethanol, dried and was used directly in the next cycle. As shown in Table 2 (entry 9-12) after the four cycle catalyst was still highly efficient.

According to the results obtained up to this point, we decided to apply this method for synthesis of mono and bis 4-amino-5-pyrimidine-carbonitrile derivatives (**4a-v**)

via three-component reaction of amidines (**1a-c**), malononitrile (**2**) and aldehydes (**3a-i**) in the presence of 30 mol% KF/Al₂O₃ in EtOH at reflux (Table 3).

CONCLUSIONS

In this study, new applications of basic, solid phase KF/Al₂O₃ for the synthesis of mono and bis 4-amino-5-pyrimidine-carbonitrile derivatives via three-component

Table 3: The three-component reaction of amidines (1a-c), malononitrile (2) and aldehydes (3a-i).

Compd. No.	R	Ar	Time (min)	Yield (%)	M. P. observed (°C)	M. P. reported (°C)
4a	NH ₂	C ₆ H ₅	5	94	287-288	288-300 [39]
4b	Ph	C ₆ H ₅	7	92	209-211	210-212 [39]
4c	NH ₂	4-CH ₃ -C ₆ H ₄	8	92	127-129	130 [39]
4d	Ph	4-CH ₃ -C ₆ H ₄	10	91	209-211	210 [39]
4e	NH ₂	4-CH ₃ O-C ₆ H ₄	10	90	237-239	240 [41]
4f	Ph	4-CH ₃ O-C ₆ H ₄	12	89	211-213	213 [39]
4g	NH ₂	4-Cl-C ₆ H ₄	4	95	227-230	229-231 [39]
4h	Ph	4-Cl-C ₆ H ₄	5	93	220-222	222 [39]
4i	NH ₂	4-Br-C ₆ H ₄	5	95	237-239	>240 (dec.) [39]
4j	Ph	4-Br-C ₆ H ₄	7	93	234-235	235-238 [39]
4k	NH ₂	Furan-2-yl	10	91	>165 (dec.)	—
4l	Ph	Furan-2-yl	12	90	173-176	—
4m	CH ₃	Furan-2-yl	12	90	>198 (dec.)	—
4n	NH ₂	Thiophen-2-yl	10	90	125-128	—
4o	Ph	Thiophen-2-yl	12	89	197-198	200 [39]
4p	CH ₃	Thiophen-2-yl	12	90	148-150	—
4q	NH ₂	Terephthalaldehyde	15	90	263 (dec.)	—
4r	Ph	Terephthalaldehyde	18	89	>300	>300 [50]
4s	CH ₃	Terephthalaldehyde	18	90	>300	>300 [50]
4t	NH ₂	Isophthalaldehyde	15	89	248-250	—
4u	Ph	Isophthalaldehyde	18	87	159-162	—
4v	CH ₃	Isophthalaldehyde	18	88	201-203	—

the reaction of amidines, malononitrile, and aldehydes are explained. KF/Al₂O₃ catalyst offers a simple, novel, and convenient method for this reaction. High yields, short reaction time, easy work-up and reusability of the catalyst are advantages of this procedure.

Acknowledgements

The authors express their great appreciation to Pharmaceutics Research Center, Institute of Neuropharmacology, Kerman University of Medical Sciences for supporting this investigation.

Received : Dec. 12, 2015 ; Accepted : Oct. 18, 2016

REFERENCES

- [1] Blass B.E., *KF/Al₂O₃ Mediated Organic Synthesis, Tetrahedron*, **58**: 9301-9320 (2002).
- [2] Yadav V.K., Kapoor K.K., *KF Adsorbed on Alumina Effectively Promotes the Epoxidation of Electron Deficient Alkenes by Anhydrous t-BuOOH, Tetrahedron*, **52**: 3659-3668 (1996).
- [3] Hu Q.S., Hu C.M., *Addition of Per(poly)fluoroalkyl Iodides to Vinyl Cyclopropane Derivatives: Synthesis of Fluoroalkyl Substituted γ,δ -Unsaturated Carbonyl Compounds, Chin. Chem. Lett.*, **8**: 565-568 (1997).

- [4] Kumar H.M.S., Reddy B.V.S., Reddy E.J., Yadav J.S., Microwave-assisted Eco-friendly Synthesis of 2-Alkylated Hydroquinones in Dry Media, *Green Chem.*, **1**: 141-142 (1999)
- [5] Rao A.K.S.B., Rao C.G., Singh D.B., Potassium Fluoride on Alumina: An Efficient Reagent for Isomerization of *N*-[2-(Alkyl/arylsulfonyl)ethyl]-2-methyl-5-nitro-1*H*-imidazoles, *Synth. Commun.*, **21**: 443-448 (1991)
- [6] Berree F., Marchand E., Morel G., Base Mediated Reactions in Solid-liquid Media: A 1,3-Dipolar Cycloaddition Route to Pyrrolines and Pyrroles from Imino Chlorosulfides, *Tetrahedron Lett.*, **33**: 6155-6158 (1992).
- [7] Boruah M., Konwar D., KF/Al₂O₃: Solid-supported Reagent Used in 1,3-Dipolar Cycloaddition Reaction of Nitrile Oxide, *Synth. Commun.*, **42**: 3261-3268 (2012).
- [8] Kharbanger I., Rohman R., Mecadon H., Myrboh B., KF-Al₂O₃ as an Efficient and Recyclable Basic Catalyst for the Synthesis of 4*H*-Pyran-3-carboxylates and 5-Acetyl-4*H*-pyrans, *International Journal of Organic Chemistry*, **2**: 282-286 (2012).
- [9] Dard B.A., Zaheer Z., Fatemaa S., Jadav S., Farooqui, M., KF-Al₂O₃ Catalyzed Domino One-pot, Three-component Synthesis of 3,5-Disubstituted-1,2,4-oxadiazoles Under Microwave-Assisted Solvent Free Conditions and Their Biological Activity, *International Journal of Pharmaceutical Research & Allied Sciences*, **4**: 93-99 (2015).
- [10] Wang X.S., Zhou J.X., Zeng Z.S., Li Y.L., Shi D.Q., Tua S.J., One-pot Synthesis of Pyrano[3,2-*c*]pyran Derivatives Catalyzed by KF/Al₂O₃, *Arkivoc*, **xi**:107-113 (2006).
- [11] Movassagh B., Soleiman-Beigi M., Nazari M., A Facile KF/Al₂O₃-mediated, One-pot Synthesis of Symmetrical Trithiocarbonates from Alkyl Halides and Carbon Disulfide, *Chem. Lett.*, **37**: 22-23 (2008).
- [12] Movassagh B., Shokri S., Potassium Fluoride Doped on Alumina: An Efficient Catalyst for Conversion of Aldoximes Into Nitriles, *Synth. Commun.*, **35**: 887-890 (2005).
- [13] Nazari M., Movassagh B., α -Phenylselenenylation of Aldehydes and Ketones with Diphenyl Diselenide Mediated by KF/Al₂O₃, *Tetrahedron Lett.*, **50**: 1453-1455 (2009).
- [14] Movassagh B., Shokri S., An Efficient and Convenient KF/Al₂O₃ Mediated Synthesis of Nitriles from Aldehydes, *Tetrahedron Lett.*, **46**: 6923-6925 (2005).
- [15] Zhu J., Bienayme H., "Multicomponent Reactions", Wiley-VCH, Weinheim (2005).
- [16] Tietze L.F., Domino Reactions in Organic Synthesis, *Chem. Rev.*, **96**: 115-136 (1996).
- [17] Tietze L.F., Beifuss U., Sequential Transformations in Organic Chemistry: A Synthesis Strategy with a Future, *Angew. Chem. Int. Ed. Engl.*, **32**: 131-163 (1993).
- [18] Mohammadi Ziarani G., Mousavi S., Lashgari N., Badii A., Shakiba M., Application of Sulfonic Acid Functionalized Nanoporous Silica (SBA-Pr-SO₃H) in the Green One-pot Synthesis of Polyhydroacridine Libraries, *Iran. J. Chem. Chem. Eng. (IJCCE)*, **32**: 9-16 (2013).
- [19] Shaabani A., Mirzaei P., Microwave-Assisted Three-Component Cyclocondensation Reaction: A Facile Synthesis of Highly Functionalized Cyclohexene Derivatives, *Iran. J. Chem. Chem. Eng. (IJCCE)*, **26**: 59-61 (2007).
- [20] Shaabani A., Bazgir A., Arab Ameri S., Sharifi Kiasaraie M., Samadi S., Comparison of Catalytic Effect of Alkali and Alkaline Earth Metals Hydrogen Sulfate: As the Promoter for an Efficient Synthesis of 3,4-Dihydropyrimidin-2(1*H*)-ones under Solvent-Free Conditions, *Iran. J. Chem. Chem. Eng. (IJCCE)*, **24**: 67-71 (2005).
- [21] Mohammadi Ziarani G., Aleali F., Lashgari N., Badii, A., An Efficient Green Approach for the Synthesis of Structurally Diversified Spirooxindoles Using Sulfonic Acid Functionalized Nanoporous Silica (SBA-Pr-SO₃H), *Iran. J. Chem. Chem. Eng. (IJCCE)*, **35**: 17-23 (2016).
- [22] Shaker R.M., The Chemistry of Mercapto- and Thione- Substituted 1,2,4-Triazoles and Their Utility in Heterocyclic Synthesis, *Arkivoc*, **ix**: 59-112 (2006).
- [23] Bellina F., Rossi R., Synthesis and Biological Activity of Pyrrole, Pyrroline and Pyrrolidine Derivatives with Two Aryl Groups on Adjacent Positions, *Tetrahedron*, **62**: 7213-7256 (2006).
- [24] Mehta P.D., Sengar N.P.S., Pathak A.K., 2-Azetidinone: A New Profile of Various Pharmacological Activities, *Eur. J. Med. Chem.*, **45**: 5541-5560 (2010).

- [25] Abaszadeh M., Seifi M., [Ultrasound-assisted 1,3-Dipolar Cycloaddition and Cyclopropanation Reactions for the Synthesis of Bis-indolizine and Bis-cyclopropane Derivatives](#), *Org. Biomol. Chem.*, **12**: 7859-7863 (2014).
- [26] Shabani M., Yousefi M., Abdouss M., Aziznejad F., Shahrosvand H., [Kinetics and Mechanism of the Substitution Reaction of Amine in \(bis\(Acetylene Acetonato\) Ethylenediimine\) \(Diamine\)Cobalt\(III\) Perchlorate with \$N_3^-\$, \$Br^-\$ and \$NCS^-\$](#) , *Iran. J. Chem. Chem. Eng. (IJCCE)*, **30**: 49-54 (2011).
- [27] Bhat A.R., Athar F., Azam A., [Bis-pyrazolines: Synthesis, Characterization and Antiamoebic Activity as Inhibitors of Growth of Entamoeba Histolytica](#), *Eur. J. Med. Chem.*, **44**: 426-431 (2009).
- [28] Iqbal P.F., Parveen H., Bhat A.R., Hayat F., Azam A., [Synthesis, Characterization, Antiamoebic Activity and Toxicity of Novel Bisdioxazole Derivatives](#), *Eur. J. Med. Chem.*, **44**: 4747-4751 (2009).
- [29] Cox R.A., [Macromolecular Structure and Properties of Ribonucleic Acids](#), *Quart. Rev. Chem. Soc.*, **22**: 499-526 (1968).
- [30] Agarwal O.P., "Organic Chemistry; Reaction and Reagent", Krishna Prakashan Media, New Delhi (2006).
- [31] Jain M.K., Sharnevsas S.C., "Modern Organic Chemistry", Jalandhar-Delhi (2008).
- [32] Mitsuya H., "Anti-HIV Nucleosides: Past, Present and Future", Chapman and Hall, New York (1997).
- [33] Kompis I., Wick A., [Synthese Von 4-Halogensubstituierten Analogen Von Trimethoprim](#), *Helv. Chim. Acta*, **60**: 3025-3034 (1977).
- [34] Hassan N.A., [Syntheses of Furo\[3,2-e\]\[1,2,4\]triazolo\[1,5-c\]pyrimidines and Furo\[2',3':5,6\]-pyrimido\[3,4-b\]\[2,3-e\]indolo\[1,2,4\] triazine as a New Ring System](#), *Molecules*, **5**: 826-834 (2000).
- [35] Cannito A., Pemmsin M., Lnu-Due C., Hoguet F., Gaultier C., Narcisse J., [Synthesis and Pharmacological Activities of Some 3-Substituted Thienopyrimidin-4-one-2-thiones](#), *Eur. J. Med. Chem.*, **25**: 635-639 (1990).
- [36] Smith P.A.S., Kan R.O., [Cyclization of Isothiocyanates as a Route to Phthalic and Homophthalic Acid Derivatives](#), *J. Org. Chem.*, **29**: 2261-2265 (1964).
- [37] Polak A., Scholer H.J., [Mode of Action of 5-Fluorocytosine and Mechanisms of Resistance](#), *Chemotherapy*, **21**: 113-130 (1975).
- [38] So H.L., Ibrahim M.E.D., [Design and Synthesis of New Anticancer Pyrimidines with Multiple-kinase Inhibitory Effect](#), *Bioorg. Med. Chem.*, **18**: 3860-3874 (2010).
- [39] Sheibani H., Saljoogi A.S., Bazgir A., [Three-component Process for the Synthesis of 4-Amino-5-pyrimidinecarbonitriles Under Thermal Aqueous Conditions or Microwave Irradiation](#), *Arkivoc*, **ii**, 115-123 (2008).
- [40] Kapupara P.P., Dholakia S.P., Patel V.P., Suhagia B.N., [Design, Synthesis and Pharmacological Screening of 4-Amino-5-pyrimidinecarbonitriles as Potential Anti-inflammatory Agents](#), *J. Chem. Pharm. Res.*, **2**: 287-294 (2010).
- [41] Deshmukh M.B., Anbhule P.V., Jadhav S.D., Jagtap S.S., Patil D.R., Salunkhe S.M., Sankpa S.A., [A Novel and Environmental Friendly, One-step Synthesis of 2,6-Diamino-4-phenyl pyrimidine-5-carbonitrile Using Potassium Carbonate in Water](#), *Indian J. Chem.*, **47B**: 792-795 (2008).
- [42] Sheibani H., Seifi M., Bazgir A., [Three-Component Synthesis of Pyrimidine and Pyrimidinone Derivatives in the Presence of High-Surface-Area MgO, a Highly Effective Heterogeneous Base Catalyst](#), *Synth. Commun.*, **39**: 1055-1064 (2009).
- [43] Ahmadi S.J., Sadjadi S., Hosseinpour M., [Granulated Copper Oxide Nanocatalyst: A Mild and Efficient Reusable Catalyst for the One-pot Synthesis of 4-Amino-5-pyrimidinecarbonitriles Under Aqueous Conditions](#), *Monatsh. Chem.*, **142**: 1163-1168 (2011).
- [44] Zahedifar M., Sheibani H., [Rapid Three-component Synthesis of Pyrimidine and Pyrimidinone Derivatives in the Presence of \$Bi\(NO_3\)_3 \cdot 5H_2O\$ as a Mild and Highly Efficient Catalyst](#), *Res. Chem. Intermed.*, **41**: 105-111 (2015).
- [45] Zare A., Hasaninejad A., Moosavi-Zare A.R., Beyzavi M.H., Khalafi-Nezhad A., Pishahang N., Parsae Z., Mahdaviniasab P., Hayati N., [\$KF/Al_2O_3\$ as a Highly Efficient Reagent for the Synthesis of N-Aryl Derivatives of Pyrimidine and Purine Nucleobases](#), *Arkivoc*, **xvi**: 178-188 (2008).

- [46] Abaszadeh M., Seifi M., [Sodium Benzenesulfonates: Novel and Effective Organo Catalyst for Three-component Synthesis 5,6,7,8-Tetrahydro-4H-chromene Derivatives Under Ultrasound Irradiation](#), *Lett. Org. Chem.*, **12**: 271-276 (2015).
- [47] Abaszadeh M., Seifi M., Asadipour A., [Ultrasound Promotes One-pot Synthesis of 1,4-Dihydropyridine and Imidazo\[1,2-a\]quinolone Derivatives, Catalyzed by ZnO Nanoparticles](#), *Res. Chem. Intermed.*, **41**: 5229-5238 (2015).
- [48] Abaszadeh M., Seifi M., [Crown Ether Complex Cation Ionic Liquids \(CECILs\) as Environmentally Benign Catalysts for Three-component Synthesis of 4,5-Dihydropyrano\[3,2-c\]chromene and 4,5-Dihydropyrano\[4,3-b\]pyran Derivatives](#), *Res. Chem. Intermed.*, **41**: 7715-7723 (2014).
- [49] Amirheidari B., Seifi M., Abaszadeh M., [Evaluation of Magnetically Recyclable Nano-Fe₃O₄ as a Green Catalyst for the Synthesis of Mono and Bis Tetrahydro-4H-chromene and Mono and Bis 1,4-Dihydropyridine Derivatives](#), *Res. Chem. Intermed.*, **42**: 3413-3423 (2016).
- [50] Shaker R.M., [Synthesis and Characterization of Some New 4,4-\(1,4-Phenylene\)dipyrimidine and 6,6-\(1,4-Phenylene\)-di\(pyridin-2\(1H\)-one\) Derivatives](#), *Heteroatom Chemistry*, **16**: 507-512 (2005).