

A Rapid Combinatorial Library Synthesis of Benzazolo[2,1-*b*]quinazolinones and Triazolo[2,1-*b*]quinazolinones

*Shaabani, Ahmad**⁺; Farhangi, Elham; Shaabani, Shabnam*

Faculty of Chemistry, Shahid Beheshti University,

P.O. Box 19396-4716 Tehran, I.R. IRAN

ABSTRACT: *Benzazolo [2,1-*b*]quinazolinones and triazolo[2,1-*b*]quinazolinones were synthesized in high yields by the condensation reaction of an aldehyde and a cyclic β -diketone with 2 - aminobenzimidazole, 2 - aminobenzothiazole or 1, 2, 4 - triazole derivatives in 1 - butyl - 3 - methylimidazolium bromide as an ionic liquid under classic heating conditions within 5-60 minutes.*

KEY WORDS: *2-Aminobenzimidazole, 2-Aminobenzothiazole, 3-Amino-1,2,4-triazole, 3,5-Diamino-1, 2,4-triazole, 3-Amino-5-thiomethoxy-1,2,4-triazole, Ionic liquid.*

INTRODUCTION

Quinazolines are reported to have a broad spectra of biological activities such as propyl hydroxylase inhibitor [1], antidiabetics [2], anti-inflammatory [3], antiviral [4], antimicrobial [5], antineoplastic [6], and potent immunosuppressive agents [7]. They also emerged as an integral backbones of calcium channel blockers[8]. So, the analogs of these heterocycles show potent antitumor activity[9,10]. Quinazoline heterocycles are the subunit of several natural alkaloids and active pharmaceuticals [1,11,12] (Scheme 1).

The useful methods, have been reported for synthesis of tetrahydrobenzimidazo [2,1-*b*] quinazolin-1(2*H*)-ones and tetrahydro-1,2,4-triazolo[5,1-*b*]quinazolin-8(4*H*)-ones ring system skeletons, are the condensation of aminoazoles with benzilidene compounds[13-16] or three-component condensation of 2-aminobenzothiazole or 2-aminobenzimidazole and an aldehyde with cyclic β -diketone [17-18]. However, these synthetic protocols

suffer from one or more disadvantages such as harsh reaction conditions, prolonged reaction times, or use of hazardous and often expensive acid catalysts. Moreover, the synthesis of these heterocycles is usually carried out in polar solvents such as methanol, acetic acid, THF and DMF leading to complex isolation and recovery procedures. These processes also generate waste containing both catalyst and solvent, which have to be recovered, treated, and disposed off.


Recently, Room-Temperature Ionic Liquids (RTILs) have attracted much attention as promising alternative 'green' solvents to hazardous traditional organic solvents, due to their properties such as non-flammability, negligible vapor pressure, high thermal stability, solvating ability and easy recyclability. They have the potential to be highly polar yet non-coordinating [19-23]. In addition to the above-mentioned salient features of Ionic Liquids (ILs) as reaction media, we have also

* To whom correspondence should be addressed.


+E-mail: a-shaabani@sbu.ac.ir

1021-9986/13/1/

12/\$/3.20



Scheme 1: Quinazoline ring containing bioactive natural products and synthetic molecules.



Scheme 2: Synthesis of tetrahydrobenzimidazo[2,1-*b*]quinazolin-1(2*H*)-ones (5*a-r*) and tetrahydro-1,2,4-triazolo[5,1-*b*]quinazolin-8(4*H*)-ones (6*a-r*)

recently shown that they can promote and catalyze important organic transformations under ambient conditions without the need for any added catalysts or ligands. The reactions investigated by us are Biginelli and Biginelli-like reaction [24,25], as well as the Groebke reaction [26], and Hantzsch reaction [27], which proceeded with significantly enhanced reaction rates and excellent isolated yields.

Due to the biological properties of quinazolinones and quinazolines and the occurrence of the quinazoline skeleton in various natural products [1,11,12,28-30] and our interest in Multi-Component Reactions (MCRs) [31-38], we developed a rapid combinatorial approach for the synthesis of tetrahydrobenzimidazo[2,1-*b*]quinazolin-1(2*H*)-ones ring systems **5** or tetrahydro-1,2,4-triazolo[5,1-*b*]quinazolin-8(4*H*)-ones ring systems **6** by a one-pot three-component condensation reaction of

an aldehyde **1** and a cyclic β -diketone **2** with 2-aminobenzimidazole, 2-aminobenzothiazole **3** or with various triazole derivatives **4** such as 3-amino-1,2,4-triazole, 3,5-diamino-1,2,4-triazole, 3-amino-5-thiomethoxy-1,2,4-triazole under conventional heating conditions without using any catalyst in 1-butyl-3-methylimidazolium bromide ([bmim]Br) (Scheme 2).

EXPERIMENTAL SECTION

Apparatus

Melting points were measured on an Electrothermal 9100 apparatus. Mass spectra were recorded on a Shimadzu QP 1100 EX mass spectrometer operating at an ionization potential of 70 eV. IR spectra were recorded on a Shimadzu IR-470 spectrometer. ¹H, and ¹³C NMR spectra were recorded on a Bruker Avance DRX-300 spectrometer at 300.13, and 75.47 MHz. NMR spectra

Table 1: Solvent effects on the reaction of dimedone and *p*-methylbenzaldehyde with 2-aminobezimidazole on the synthesis of 3,3-dimethyl-12-(4-methyl-phenyl)-3,4,5,12-tetrahydrobenzimidazo[2,1-*b*]quinazolin-1(2*H*)-one 5b^a.

Entry	Solvent	Time (h)	Temperature (°C)	Yield (%)
1	CH ₂ Cl ₂	2	Reflux	0
2	CHCl ₃	2	Reflux	0
3	CH ₃ COOEt	2	Reflux	0
4	CH ₃ C ₆ H ₅	2	Reflux	0
5	CH ₃ CN	2	Reflux	25
6	CH ₃ OH	2	Reflux	43
7	CH ₃ CH ₂ OH	2	Reflux	66
8	Water	6	80	trace
9	[Bmim]Br	2	80	81
10	[Bmim]Br	1	80	79
11	[Bmim]Br	0.5	80	79
12	-	2	80	trace

a) 2-Aminobenzimidazole (1 mmol), dimedone (1 mmol) and *p*-methylbenzaldehyde (3 mmol).

were obtained on solutions in DMSO-d⁶. The elemental analysis was performed with an Elementar Analysensysteme GmbH VarioEL CHNS mode. All the products are known compounds, and were characterized by melting points, IR, ¹H NMR, ¹³C NMR and mass spectra. In some cases we used CHNS for compounds which because of low solubility we could not obtain ¹³C NMR.

General procedure for synthesis of 3,4,5,12-tetrahydrobenzimidazo [2,1-*b*]quinazolin-1(2*H*)-one (5a-r)

A mixture of cyclic β-diketones (1 mmol), aldehydes (3 mmol), and 2-aminobenzazoles (1 mmol) was successively added to a screw-capped vial containing [bmim]Br (0.3 g) and a magnetic stirring bar. The mixture was heated at 100 °C in a preheated oil bath for 5-60 min. Then, the reaction mixture was washed with cold water and filtered. The solid residue was washed with acetone (3×5 mL) to produce **5** as a white powder in high yield.

General procedure for synthesis of tetrahydro-1,2,4-triazolo[5,1-*b*]quinazolin-8(4*H*)-ones(6a-r)

A mixture of cyclic β-diketones (1 mmol), aldehydes (1 mmol), and amino-1,2,4-triazoles (1 mmol) was added to a screw-capped vial containing [bmim]Br (0.3 g) and a magnetic stirring bar. The reaction mixture was heated

at 80 °C in a preheated oil bath for 15-30 min. Then, the reaction mixture was washed with cold water and filtered. The solid residue was washed with acetone (3×5 mL) to produce **6a-r** as a white powder in high yields.

RESULTS AND DISCUSSION


The one-pot method involves the classical heating of an aldehyde **1** and a cyclic β-diketone **2** with 2-aminobenzothiazole or 2-aminobezimidazole **3** or 3-amino-1,2,4-triazole or 3,5-diamino-1,2,4-triazole or 3-amino-5-thiomethoxy-1,2,4-triazole **4** without using any catalyst in [bmim]Br to give a family of tetraheterocyclic tetrahydrobenzimidazo[2,1-*b*]quinazolin-1(2*H*)-ones **5** and triheterocyclic tetrahydro-1,2,4-triazolo[5,1-*b*]quinazolin-8(4*H*)-ones **6** in high yields.

To achieve suitable conditions for the above transformations, series of experiments were carried out (Tables 1 and 2). First we investigated the reaction of dimedone and *p*-methylbenzaldehyde with 2-aminobezimidazole under solvent-free conditions, various solvents, water and ionic liquids under classic heating conditions. In the absence of solvent, the reaction was very slow and the yield of the product was trace at 80 °C after 2 h (Table 1, Entry 12). The reaction did not proceed in aprotic organic solvents even under reflux conditions after 2 h (Table 1, Entries 1-5). The yield of product was improved in protic organic solvents (Table 1, Entries 6-7), but we

Table 2: One-pot synthesis of tetrahydrobenzimidazo[2,1-b]quinazolin-1(2H)-one 5b at different ionic liquid at 30 min^a.

Entry	Ionic liquid	Temperature (°C)	Yield (%)
1	BPB	80	48
2	TMGT	80	43
3	TMGA	80	57
4	TBAB	80	81
5	TBAC	80	79
6	MIT	80	64
7	[bmim]Br	80	89
8	[bmim]Br	100	96
9	[bmim]Br	120	96
10	[bmim]PF ₆	80	82
11	-	80	trace

a) 2-Aminobenzimidazole (1 mmol), dimedon (1 mmol) and p-methylbenzaldehyde (3 mmol).

**Scheme 3: Reaction optimization for the synthesis of 5b**

found the reaction yield was trace in water after 6 h at 80 °C (Table 1, Entry 8). The best solvent was found to be TetraButyl Ammonium Bromide (TBAB) and the reaction yield was 79% after heating 30 min at 80°C (Table 1, Entry 11).

In order to examine the best ionic liquid, different ionic liquids such as Butyl Pyridinium Bromide (BPB), Tetra Methyl Guanidinium Trifluoroacetate (TMGT), Tetra Methyl Guanidinium Acetate (TMGA), TetraButyl Ammonium Bromide (TBAB), TetraButyl Ammonium Chloride (TBAC), Methylimidazolium Trifluoroacetate (MIT), 1-butyl-3-methylimidazolium bromide ([bmim]Br), 1-butyl-3-methylimidazolium hexafluorophosphate ([bmim]PF₆) in this condensation reaction were applied (Table 2). In the course of this study it was found that 1-butyl-3-methylimidazolium bromide was the best ionic liquid for this reaction in terms of yield, price and easy at work-up.

To illustrate the need for [bmim]Br in these reactions, the reaction of aldehyde **1**, cyclic β -diketone **2** and 2-aminobenzimidazole **3** was studied in the absence of [bmim]Br. The yield of product was trace at 80 °C after 2 h (Table 2, Entry 11).

In order to improve the yields, we performed reaction using different quantities of reagents. The best result was obtained with 3:1:1 molar ratios of aldehyde, cyclic β -diketone and 2-aminobenzimidazole or 2-aminobenzothiazole. However, the 1:1:1 molar ratios of these reagents gave the compound **7** (Scheme 3). In the case of 3-amino-1,2,4-triazole or 3,5-diamino-1,2,4-triazole or 3-amino-5-thiomethoxy-1,2,4-triazole, the optimum molar ratio was 1:1:1.

One of the advantages of Ionic Liquids (ILs) is the ability to be recyclable as reaction medium. We were able to separate [bmim]Br from reaction medium easily by washing with water and evaporating the solvent

Table 3: One-pot synthesis of tetraheterocyclic tetrahydrobenzimidazo[2,1-b]quinazolin-1(2H)-ones by the condensation reaction of an aldehyde, a cyclic β -diketone and 2-aminobenzothiazole or 2-aminobenzimidazole in [bmim]Br at 100 °C after 5-60 min.

Product	Y	Z	X	Yield (%)	Time(min)
5a	NH	C(CH ₃) ₂	H	94	5
5b	NH	C(CH ₃) ₂	4-CH ₃	96(94,93,92,93) ^a	10
5c	NH	C(CH ₃) ₂	4-OCH ₃	89	15
5d	NH	C(CH ₃) ₂	4-Cl	91	10
5e	NH	C(CH ₃) ₂	4-Br	93	10
5f	NH	C(CH ₃) ₂	3-OCH ₃	85	15
5g	NH	C(CH ₃) ₂	3-Cl	89	5
5h	NH	C(CH ₃) ₂	3-Br	91	15
5i	NH	C(CH ₃) ₂	3-NO ₂	88	15
5j	NH	C(CH ₃) ₂	2-CH ₃	94	5
5k	NH	C(CH ₃) ₂	2-Cl	91	10
5l	NH	CH ₂	H	86	20
5m	NH	CH ₂	4-CH ₃	91	15
5n	NH	CH ₂	4-OCH ₃	88	20
5o	NH	CH ₂	3-NO ₂	90	15
5p	S	C(CH ₃) ₂	H	73	60
5q	S	C(CH ₃) ₂	2-OCH ₃	69	60
5r	S	C(CH ₃) ₂	3-OCH ₃	72	60

a) The same ionic liquid runs for four times.

under vacuum, and reuse it for subsequent reactions (Table 3, Product **5b**).

To explore the scope and limitations of this reaction, we have extended it to various *ortho*-, *meta*- and *para*-substituted benzaldehydes in the presence of 2-aminobenzimidazole, 2-aminobenzothiazole, 3-amino-

1,2,4-triazole and 3,5-diamino-1,2,4-triazole or 3-amino-5-thiomethoxy-1,2,4-triazole. As indicated in Tables 3 and 4, the reaction proceeded efficiently with benzaldehyde and electron-withdrawing and electron-donating *ortho*-, *meta*- and *para*-substituted benzaldehydes.

Table 4: One-pot synthesis of tetrahydro-1,2,4-triazolo[5,1-b]quinazolin-8(4H)-ones by the condensation reaction of an aldehyde, a cyclic β -diketone and 3-amino-1,2,4-triazole, 3,5-diamino-1,2,4-triazole, 3-amino-5-thiomethoxy-1,2,4-triazole in [bmim]Br at 80 °C after 15-30 min.

Product	Y	Z	X	Yield (%)	Time(min)
6a	H	C(CH ₃) ₂	H	93	20
6b	H	C(CH ₃) ₂	4-CH ₃	96	15
6c	H	C(CH ₃) ₂	4-OCH ₃	86	20
6d	H	C(CH ₃) ₂	4-Cl	87	20
6e	H	C(CH ₃) ₂	3-OCH ₃	89	20
6f	H	C(CH ₃) ₂	3-NO ₂	91	30
6g	H	C(CH ₃) ₂	2-CH ₃	93	20
6h	H	CH ₂	H	90	30
6i	H	CH ₂	4-CH ₃	94	20
6j	H	CH ₂	4-OCH ₃	85	20
6k	H	CH ₂	4-Cl	87	30
6l	H	CH ₂	3-NO ₂	89	30
6m	NH ₂	C(CH ₃) ₂	H	93	20
6n	NH ₂	C(CH ₃) ₂	4-CH ₃	87	20
6o	NH ₂	C(CH ₃) ₂	4-OCH ₃	84	20
6p	NH ₂	C(CH ₃) ₂	4-Cl	90	30
6q	SCH ₃	C(CH ₃) ₂	H	85	20
6r	SCH ₃	C(CH ₃) ₂	4-OCH ₃	84	20

CONCLUSIONS

Three - component condensation of 2-aminobenzimidazole, 2-aminobenzothiazole or various 3-amino-1,2,4-triazoles, aldehydes, and 1,3-diones in ionic liquid afforded the novel library of quinazoline molecules. It has been demonstrated that the ionic liquids are very effective in speeding up both the soluble supported multistep synthesis, as well as accelerating the rate of subsequent multicomponent reaction. Great substrate scope, high yields, short reaction time, pure products and easy work up are advantages of this procedure

in comparison to other methods.

This approach provides a high speed path for the rapid synthesis of molecular libraries with high degree of structural diversity.

Acknowledgement

Financial assistance from the Research Council of Shahid Beheshti University of Iran is gratefully acknowledged.

Received : Jun. 5, 2012 ; Accepted : Aug. 14, 2012

REFERENCES

- [1] Chai D., Fitch D. M., Propyl Hydroxylase Inhibitors. WO Patent 09039322 A1, March (2009).
- [2] Kato F., Kimura H., Omatsu M., Yamamoto K., Miyamoto R., Preventive or Therapeutic Medicines for Diabetes Containing Fused-Heterocycle Compounds or Their Salts. WO Patent 02040485, May 23 (2002).
- [3] McMaster B., Anti-Inflammatory Compositions and Methods of Use. WO Patent 03105857, December 24 (2003).
- [4] Shigeta S., Mori S., Baba M., Hosoya M., Mochizuki N., Chiba T., De Clercq E., Inhibitory Effect of Pyridobenzoazoles on Orthomyxo- and Paramyxovirus Replication in Vitro, *Antiviral Chem. Chemother.*, **3**, p. 171 (1992).
- [5] Nofal Z. M., Fahmy H. H., Mohamed, H. S., Synthesis, Antimicrobial and Molluscicidal Activities of New Benzimidazole Derivatives, *Arch. Pharm. Res.*, **25**, p. 28 (2002).
- [6] Hafez A., Monem A. A., Benzimidazole Condensed Ring Systems: New Synthesis and Antineoplastic Activity of Substituted 3,4-dihydro- and 1,2,3,4-Tetrahydro-benzo[4,5]imidazo[1,2-a]pyrimidine Derivatives, *Arch. Pharm. Res.*, **30**, p. 678 (2007).
- [7] Lunn W. H. W., Harper R. W., Stone R. L., Benzimidazo[2,1-b]quinazolin-12-ones. A New Class of Potent Immunosuppressive Compounds, *J. Med. Chem.*, **14**, p. 1069 (1971).
- [8] Alajarin R., Vaquero J. J., Alvarez-Builla J., Faude Casa-Juana M., Sunkel C., Priego J. G., Gomez-Sal P., Torres R., Imidazo[1,5-a]pyrimidine and Benzo[4,5]imidazo[1,2-a]pyrimidine Derivatives as Calcium Antagonists, *Bioorg. Med. Chem.*, **2**, p. 323 (1994).
- [9] Ishida J., Wang H. K., Bastow K. F., Hu C. Q., Lee K. H., Antitumor Agents 201. Cytotoxicity of Harmine and β -carboline Analogs, *Bioorg. Med. Chem. Lett.*, **9**, p. 3319 (1999).
- [10] Da Settimo A., Da Settimo F., Marini A. M., Primofiore G., Salerno S., Viola G., Dalla Via L., Magno S. M., Synthesis, DNA Binding and in Vitro Antiproliferative Activity of Purinoquinazoline, Pyridopyrimidopurine and Pyridopyrimidobenzimidazole Derivatives as Potential Antitumor Agents, *Eur. J. Med. Chem.*, **33**, p. 685 (1998).
- [11] Wu Y. C., Duh C. Y., Two New Natural Azafluorene Alkaloids and a Cytotoxic Aporphine Alkaloid from *Polyalthialongifolia*, *J. Nat. Prod.*, **53**, p. 1327 (1990).
- [12] He X. F., Wang X. N., Fan C. Q., Gan L. S., Yin S., Yue J. M., Chemical Constituents of *Polyalthianemorialis*, *Helv. Chim. Acta.*, **90**, p. 783 (2007).
- [13] Wendelin W., Gößnitzer E., Abou El Ella D., Synthesis and Structure Elucidation of Pyrimidobenzimidazoles and Fused Derivatives II [1]. Dodecahydrobenzimidazo-[2,1-b]quinazolines and Decahydrobenzimidazo-[2,1-b]benzo [h] quinazolines[2], *Monatsh. Chem.*, **131**, p. 353 (2000).
- [14] El Ella D. A., Gößnitzer E., Wendelin W., Synthesis and Structural Elucidation of Pyrimido-[1,2-a]benzimidazoles and Fused Derivatives. I. Dihydropyrimido[1,2-a]benzimidazoles, *J. Heterocyclic Chem.*, **33**, p. 373 (1996).
- [15] Lipson V. V., Desenko S. M., Ignatenko I. V., Shishkin O. V., Shishkina S. V., Synthesis and Chemical Transformations of Partially Hydrogenated [1,2,4]triazolo[5,1-b]quinazolines, *Russ. Chem. Bull.*, **55**, p. 345 (2006).
- [16] Shikhaliev Kh. S., Kryl'skii D. V., Yu. Potapov A., Krysin M. Y., Synthesis of 8,8-R,8,9-dihydro[1,2,4]triazolo[1,5-a]quinazolin-6(7H)-ones, *Russ. Chem. Bull.*, **54**, p. 2903 (2005).
- [17] Shaabani A., Farhangi E., Rahmati A., Synthesis of Tetrahydrobenzimidazo[1,2-b]quinazolin-1(2H)-one and tetrahydro-1,2,4-triazolo[5,1-b]quinazolin-8(4H)-one Ring Systems under Solvent-free Conditions, *Comb. Chem. High Throughput Screening*, **9**, p. 771 (2006).
- [18] Lipson V. V., Desenko S. M., Borodina V. V., Shirobokova M. G., Musatov V. I., Synthesis of Partially Hydrogenated 1,2,4-triazoloquinazolines by Condensation of 3,5-diamino-1,2,4-triazole with Aromatic Aldehydes and Dimedone, *Russ. J. Org. Chem.*, **41**, p. 114 (2005).
- [19] Wasserscheid P., Welton T., "Ionic Liquids in Synthesis", Wiley-VCH, (2003).
- [20] Wasserscheid P., Keim W., Ionic liquids - New Solutions for Transition Metal Catalysis, *Angew. Chem., Int. Ed.*, **39**, p. 3773 (2000).
- [21] Cole A. C., Jensen J. L., Ntai I., Loan K., Tran K. L., Weaver T. K. J., Forbes D. C., Jr. Davis J. H., Novel Brønsted Acidic Ionic Liquids and Their use as Dual Solvent-catalysts, *J. Am. Chem. Soc.*, **124**, p. 5962 (2002).

- [22] Welton T., Ionic liquids in catalysis, *Cordin. Chem. Rev.*, **248**, p. 2459 (2004).
- [23] Jain N., Kumar A., Chauhan S., Chauhan S. M. S., Chemical and Biochemical Transformations in Ionic Liquids, *Tetrahedron*, **61**, p. 1015(2005).
- [24] Shaabani A., Rahmati A., Ionic Liquid Promoted Efficient Synthesis of 3,4-dihydropyrimidin-2-(1H)-ones, *Catal. Lett.*, **100**, p. 177 (2005).
- [25] Shaabani A., Rahmati A., Naderi S., A Novel one-pot Three-component Reaction: Synthesis of Triheterocyclic 4H-pyrimido[2,1-b]benzazoles Ring Systems, *Bioorg. Med. Chem. Lett.*, **15**, p. 5553 (2005).
- [26] Shaabani A., Soleimani E., Maleki A., Ionic Liquid Promoted One-pot Synthesis of 3-Aminoimidazo[1,2-a]pyridines, *Tetrahedron Lett.*, **47**, p. 3031 (2006).
- [27] Shaabani A., Rezayan A. H., Rahmati A, Sharifi M., Ultrasound-Accelerated Synthesis of 1,4-dihydropyridines in an Ionic Liquid, *Monatsh. Chem.*, **137**, p. 77 (2006).
- [28] Chapman N. B., Gibson G. M., Mann F. G., Synthetic Antimalarials. Part XVI. 4-Dialkylaminoalkylaminoquinazolines. Variation of Substituents in the 6- and 7-positions, *J. Chem. Soc.*, p. 890 (1947).
- [29] Liéby-Muller F., Simon C., Constantieux T., Rodriguez J., Current developments in Michael Addition-based Multicomponent Domino Reactions Involving 1,3-dicarbonyls and Derivatives, *QSAR Comb. Sci.*, **25**, p. 432 (2006).
- [30] Simon C., Constantieux T., Rodriguez J., Utilisation of 1,3-dicarbonyl Derivatives in Multicomponent Reactions, *Eur. J. Org. Chem.*, p. 4957 (2004).
- [31] Shaabani A., Maleki A., Moghimi-Rad J., A Novel Isocyanide-based Three-component Reaction: Synthesis of Highly Substituted 1,6-dihydropyrazine-2,3-dicarbonitrile Derivatives, *J. Org. Chem.*, **72**, p. 6309 (2007).
- [32] Shaabani A., Maleki A., Mofakham H., Moghimi-Rad J., A Novel One-pot Pseudo-five-component Synthesis of 4,5,6,7-tetrahydro-1H-1, 4-diazepine-5-carboxamide Derivatives, *J. Org. Chem.*, **73**, p. 3925 (2008).
- [33] Shaabani A., Maleki A., Mofakham H., Khavasi H. R., Novel Isocyanide-based Three-component Synthesis of 3,4-dihydroquinoxalin- 2-amine derivatives, *J. Comb. Chem.*, **10**, p. 323 (2008).
- [34] Shaabani A., Maleki A., Mofakham H., Novel Multicomponent One-pot Synthesis of Tetrahydro-1H-1,5-benzodiazepine- 2-carboxamide Derivatives, *J. Comb. Chem.*, **10**, p. 595 (2008).
- [35] Shaabani A., Maleki A., Hajishaabanha F., Mofakham H., Seyyedhamzeh M., Mahyari M., Ng S. W., Novel Syntheses of Tetrahydrobenzodiazepines and Dihydropyrazines via Isocyanide-based Multicomponent Reactions of Diamines, *J. Comb. Chem.*, **12**, p. 186 (2010).
- [36] Shaabani A., Rezayan A. H., Sarvary A., Keshipour S., Ng S. W., A Novel One-pot Three-(in situ Five-) Component Condensation Reaction: An Unexpected Approach for the Synthesis of Tetrahydro-2,4-dioxo-1H-benzo[b][1,5] diazepine-3-yl-2-methylpropanamide Derivatives, *Org. Lett.*, **11**, p. 3342 (2009).
- [37] Shaabani A., Soleimani E., Rezayan A. H., Sarvary A., Khavasi H. R., Novel Isocyanide-based Four-Component Reaction: A Facile Synthesis of Fully Substituted 3,4-dihydrocoumarin Derivatives, *Org. Lett.*, **10**, p. 2581 (2008).
- [38] Shaabani, A.; Mofakham, H.; Maleki, A.; Hajishaabanha, F., Novel Isocyanide-based One-pot Multicomponent Syntheses of Tetrahydrobenzo[b] [1,4]oxazepine and Malonamide Derivatives, *J. Comb. Chem.*, **12**, p. 630 (2010).