

# One Pot Synthesis of Polyhydroquinolines Catalyzed by Sulfonic Acid Functionalized SBA-15 as a New Nanoporous Acid Catalyst under Solvent Free Conditions

**Mohammadi Ziarani, Ghodsi\*<sup>+</sup>**

*Department of Chemistry, University of Alzahra, Tehran, I.R. IRAN*

**Badiei, Ali Reza; Khaniania, Yeganeh**

*Faculty of Chemistry, University of Tehran, Tehran, I.R. IRAN*

**Haddadpour, Mahbobeh**

*Department of Chemistry, University of Alzahra, Tehran, I.R. IRAN*

**ABSTRACT:** Sulfonic acid functionalized SBA-15 (SBA-Pr-SO<sub>3</sub>H) as a new nanoporous acid catalyst was used in the one-pot synthesis of polyhydroquinolines derivatives via the Hantzsch four component condensation reaction of aldehydes,  $\beta$ -ketoesters, dimedone and ammonium acetate under solvent free conditions with short reaction time in excellent yields. SBA-Pr-SO<sub>3</sub>H was proved to be an efficient heterogeneous nanoporous solid acid catalyst (pore size 6 nm), which could be easily handled and removed from the reaction mixture by simple filtration, and also recovered and reused without loss of reactivity.

**KEY WORDS:** Nanoporous solid acid catalyst, Sulfonic acid functionalized SBA-15, Polyhydroquinolines, Solvent free conditions, Dimedone.

## INTRODUCTION

1,4-Dihydropyridines possess a variety of biological activities, such as vasodilator, bronchodilator, antiatherosclerotic, antitumor, geroprotective, hepatoprotective, and antidiabetic agents [1-2].

Several efficient conditions and methods have been developed for the synthesis of 1,4-dihydropyridines such as

*Hantzsch* method [3], the use of microwave [4], ionic liquid [5], high temperature in refluxing solvent [6], TMSCl-NaI [7], metal triflates Yb(OTf)<sub>3</sub> [8], molecular iodine [9] and organocatalysts [10]. However, the use of high temperatures, expensive metal precursors and long reaction times are limiting these methods. Thus,

---

\* To whom correspondence should be addressed.

+ E-mail: gmziarani@gmail.com

1021-9986/10/2/1

10/\$/3.00

the development of a simple, efficient and versatile method for the preparation of 1,4-dihydropyridine derivatives is great tendency on this research area and the entering of further improvement towards the milder reaction conditions and higher product yields is of importance in this process.

The high ordered nanoporous silica, such as MCM-41 [10], LUS-1 [11, 12] and SBA-15 [13] are unique inorganic solid supports that have very high surface area with controllable pore sizes between 2 to 30 nm. They can be employed as catalysts [14, 15], for the preconcentration of metals [16-18], and as modified carbon electrodes [19-21]. The SBA-15 is new nanoporous silica with hexagonal structure, large pore, high surface area, high thermal stability and also diffusion free due to thicker pore walls and larger pore size respectively. This can be prepared by using commercially available triblock copolymer pluronic P126 as a structure directing agent [13]. Integration of acidic functional groups (e.g.,  $-\text{SO}_3\text{H}$ ) into SBA-15 has been explored to produce promising solid acids. The sulfonic acid functionalized SBA-15 were usually synthesized through direct synthesis or post-grafting [22-23]. There have been only a few reports about the application of several types of sulfonic acid functionalized ordered mesoporous silicas as nano acid catalyst in chemical transformations [24]. For example, SBA-Pr- $\text{SO}_3\text{H}$  has been used in the synthesis of chromenes from chromanols [25], and the *von Pechmann* reaction [26].

In this work, we would like to explore the catalytic activity of the Sulfonic acid functionalized SBA-15 (SBA-Pr- $\text{SO}_3\text{H}$ ) as a highly efficient heterogenous nanoporous acid catalyst towards the one-pot synthesis of polyhydroquinolines.

## EXPERIMENTAL SECTION

### Apparatus

Gc-Mass analysis was performed on a Gc-Mass model: 5973 network mass selective detector, Gc 6890 Agilent. IR spectra were recorded from KBr disk using a FT-IR Bruker Tensor 27 instrument. Melting points were measured by using the capillary tube method with an electro thermal 9200 apparatus. The  $^1\text{H-NMR}$  (250 MHz) was run on a Bruker DPX, 250 MHz. Nitrogen adsorption and desorption isotherms were measured at  $-196^\circ\text{C}$  using a Japan Belsorb II system after the samples were vacuum dried at  $150^\circ\text{C}$  overnight. Surface areas

were calculated by the Brunauer-Emmett-Teller (BET) method, and pore sizes were calculated by the Barrett-Joyner-Halenda (BJH) method. Thermogravimetry analysis (TGA) was carried out in Perkin Elmer Pyris Diamond instrument from ambient temperature to  $800^\circ\text{C}$  using  $20^\circ\text{C}/\text{min}$  ramp rate.

### Preparation of SBA-15


At First, pluronic P123 triblock copolymer surfactant ( $\text{EO}_{20}\text{PO}_{70}\text{EO}_{20}$ ,  $M_{ac}=5800$ ) (4.0 g) was dissolved in 30 g of water and 120 g of 2 M HCl solution. Then, TEOS (tetraethyl orthosilicate) (8.50 g) was added to reaction mixture which was stirred for 8 h at  $40^\circ\text{C}$ . The resulting mixture was transferred into a teflon-lined stainless steel autoclave and kept at  $100^\circ\text{C}$  for 20 h without stirring. The gel composition P123: HCl:  $\text{H}_2\text{O}$ : TEOS was 0.0168:5.854: 162.681: 1 in molar ratio. After cooling down to room temperature, the product was filtered, washed with distilled water and dried overnight at  $60^\circ\text{C}$  in air. The as-synthesized sample was calcinated at  $550^\circ\text{C}$  for 6 h in air atmosphere to remove the copolymer template.

### Functionalization of the SBA-15 by organic groups

Functionalization of the SBA-15 catalyst was performed according to Fig. 3. The calcinated SBA-15 (2 g) and (3-mercaptopropyl)trimethoxysilane (10 mL) in dry toluene (20 mL) were refluxed for 24 h. The product was filtered and extracted for 6h in  $\text{CH}_2\text{Cl}_2$  using a soxhlet apparatus, then dried under vacuum. The solid product was oxidized with  $\text{H}_2\text{O}_2$  (excess) and one drop of  $\text{H}_2\text{SO}_4$  in methanol (20 mL) for 24 h at rt and then the mixture was filtered and washed with  $\text{H}_2\text{O}$ , and acetone. The modified SBA-15-Pr- $\text{SO}_3\text{H}$  was dried and used as nanoporous solid acid catalyst in the following reaction.

### General method for the preparation of polyhydroquinolines derivatives

The SBA-Pr- $\text{SO}_3\text{H}$  (0.05 g) was activated in vacuum at  $100^\circ\text{C}$  and then after cooling to room temperature, aldehyde 1 (1.0 mmol), 5,5-dimethyl-1,3-cyclohexanedione 2 (1.0 mmol), ethyl acetoacetate 3 (1.0 mmol), ammonium acetate (1.5 mmol) were added to catalyst. Then the reaction mixture was stirred at  $90^\circ\text{C}$  for 5-15 min and a solid product formed. After the completion of reaction as indicated by TLC, the resulting solid product was dissolved



**Fig. 1:** Synthesis of polyhydroquinoline derivatives from aldehyde, dimedone,  $\beta$ -ketoesters, and ammonium acetate in the presence of SBA-Pr-SO<sub>3</sub>H as efficient nano acid catalyst.

in hot ethanol, filtered for removing the unsolvable catalyst and then the filtrate was cooled to afford the pure product as a yellow solid. The spectroscopic and analytical data for selected compounds are presented in the following part. The catalyst was washed subsequently with acetonitrile, diluted acid solution, distilled water and then acetone, dried under vacuum and re-used for several times without loss of significant activity.

**2, 7, 7-Trimethyl - 5 - oxo-4-(2,3 - dichlorophenyl)-1,4,5, 6,7, 8 - hexa-hydroquinoline - 3 - carboxylic acid ethyl ester 4b**

Yellow solid; m.p: 229-234 °C; IR(KBr): 3290, 3074, 2959, 1700, 1642, 1605, 1484 cm<sup>-1</sup>. <sup>1</sup>H NMR (250 MHz, CDCl<sub>3</sub>):  $\delta$  = 0.96 (s, 3H, CH<sub>3</sub>), 1.07 (s, 3H, CH<sub>3</sub>), 1.15 (t, 3H, J=7Hz, CH<sub>2</sub>CH<sub>3</sub>), 2.14-2.33 (m, 4H, 2\*CH<sub>2</sub>), 2.35 (s, 3H, CH<sub>3</sub>), 4.02 (q, 3H, J=7Hz, CH<sub>2</sub>CH<sub>3</sub>), 5.43 (s, 1H, CH), 5.76 (s, 1H, NH), 7.05 (t, 1H, J=8Hz, ArH), 7.21 (dd, 1H, ArH), 7.31 (dd, 1H, ArH). MS: m/z (%) 407 [M<sup>+</sup>], 372, 262, 234.

**2,7,7-Trimethyl -5- oxo-4-Isopropyl-1,4,5,6,7,8 - hexa- hydroquinoline -3-carboxylic acid ethyl ester 4e**

Yellow solid; m.p: 162-164 °C; IR(KBr): 3279, 3209, 3076, 2958, 1679, 1601, 1486, 1386, 1331, 1222 cm<sup>-1</sup>. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>):  $\delta$  = 1.00 (s, 6H, 2\*CH<sub>3</sub>), 1.08 (s, 6H, 2\*CH<sub>3</sub>), 1.23 (t, 3H, J = 7.2 Hz, CH<sub>2</sub>H<sub>3</sub>), 2.07-2.21 (m, 6H), 2.25 (s, 3H, CH<sub>3</sub>), 4.01 (q, 2H, J = 7.2 Hz, CH<sub>2</sub>CH<sub>3</sub>), 5.86 (s, 1H, NH). Ms: m/z (%) 304[M<sup>+</sup>], 262, 234, 218, 205, 178.

**2,7,7- Trimethyl-5-oxo-4- (3,4-dimethoxyphenyl)-1,4,5,6,7,8 - hexa- hydroquinoline- 3- carboxylic acid ethyl ester 4l**

Yellow solid; m.p: 207-209 °C; IR (KBr): 3278, 3078, 2936, 1965, 1603, 1489, 1308, 1217 cm<sup>-1</sup>. <sup>1</sup>H NMR (250

MHz, CDCl<sub>3</sub>):  $\delta$  = 0.95 (s, 3H, CH<sub>3</sub>), 1.08 (s, 3H, CH<sub>3</sub>), 1.21 (t, 3H, J = 7Hz, CH<sub>2</sub>CH<sub>3</sub>), 2.2-2.32 (m, 4H, 2\*CH<sub>2</sub>), 2.38 (s, 3H, CH<sub>3</sub>), 3.80 (s, 3H, OCH<sub>3</sub>), 3.83 (s, 3H, OCH<sub>3</sub>), 4.08 (q, 2H, J= 7 Hz, CH<sub>2</sub>CH<sub>3</sub>), 5.01 (s, 1H, CH), 5.63 (s, 1H, NH), 6.68-6.76 (m, 2H, ArH), 6.92 (d, 1H, ArH). Ms: m/z (%) 399 [M<sup>+</sup>] 370, 354, 326, 262, 234.

**2,7,7-Trimethyl-5-oxo-4-(3-methoxyphenyl)-1,4,5, 6, 7, 8 - hexa- hydroquinoline-3-carboxylic acid ethyl ester 4m**

Yellow solid, m.p: 208-211. IR (KBr): 3302, 3076, 2959, 1696, 1645, 1610, 1483 cm<sup>-1</sup>. <sup>1</sup>H NMR (250 MHz, CDCl<sub>3</sub>):  $\delta$  = 0.95 (s, 3H, CH<sub>3</sub>), 1.07 (s, 3H, CH<sub>3</sub>), 1.2 (t, 3H, J=7 Hz, CH<sub>2</sub>CH<sub>3</sub>), 2.12-2.28 (m, 4H, 2\*CH<sub>2</sub>), 2.39 (s, 3H, CH<sub>3</sub>), 3.75 (s, 3H, OCH<sub>3</sub>), 4.08 (q, 2H, J=7 Hz, CH<sub>2</sub>CH<sub>3</sub>), 5.04 (s, 1H, CH), 5.93 (s, 1H, NH), 6.62-7.2 (m, 5H, Ar H). Ms: m/z (%) 369 [M<sup>+</sup>], 340, 324, 296, 262, 234.

**RESULTS AND DISCUSSION**

The classical method for the preparation of polyhydroquinolines derivatives involves the reaction of aldehydes with ethyl acetoacetate, and ammonia in acetic acid or in refluxing alcohol. However, this method suffers from several disadvantages such as long reaction times, excess of organic solvent, lower product yields and refluxing conditions. Therefore, after using MCM-41 in this synthesis by Nagarapu [27], and in continuation of our studies on the application of new acid catalysts in organic synthesis [28], we had the opportunity to explore the catalytic activity of nanoporous catalyst of SBA-Pr-SO<sub>3</sub>H towards the synthesis of 1,4-dihydropyridines from aldehyde, dimedone,  $\beta$ -ketoesters, and ammonium acetate under solvent free conditions as shown in Fig. 1. The reaction has efficiently been carried out in excellent yields with short reaction time. The results were summarized in Table 1.

Table 1: The SBA-Pr-SO<sub>3</sub>H catalyzed the synthesis of polyhydroquinoline derivatives.





























Entry	Aldehyde	Product	Time (min)	Yield (%)	mp (°C)	Ref.
1		 4a	13	80	241-244	241-244 [29b]
2		 4b	13	80	229-234	--
3		 4c	12	90	204-206	204-206 [30]
4		 4d	12	85	231-233	229-231 [29b]
5		 4e	18	80	162-166	-
6		 4f	10	85	227-230	227-229 [31]
7		 4g	10	90	260-262	260-262 [31]

Table 1: (continued)

8			12	90	264-266	260-261 [32]
9			12	85	242-244	245-246 [29a]
10			12	90	239-240	238-240 [32]
11			10	85	242-244	242-244 [30]
12			10	90	207-209	-
13			10	90	208-211	--
14			14	75	241-244	241-244 [29b]




Fig. 2: The proposed mechanism.




Fig. 3: Schematic illustration for the preparation of SBA-Pr-SO<sub>3</sub>H.

The proposed mechanism was shown in Fig. 2. The polyhydroquinoline derivatives could be synthesized by two methods. The SBA-Pr-SO<sub>3</sub>H catalyses subsequently the *Knoevenagel* type coupling of aldehydes with active methylene compounds and then the *Michael* type addition of intermediates together to provide the products [29a].

### ***Knoevenagel condensation***

#### *Preparation and characterization of catalyst*

Pure Nanoporous compound SBA-15 was synthesized according to the well-established method designed by Zhao & coworkers [13] with triblock poly(ethylene oxide)-b-poly(propylene oxide)-b-poly(ethylene oxide) copolymer (Pluronic, EO<sub>20</sub>PO<sub>70</sub>EO<sub>20</sub>, P123) as the template. A schematic illustration for the preparation of SBA-Pr-SO<sub>3</sub>H was shown in Fig. 3. First, the calcined SBA-15 silica was functionalized with (3-mercaptopropyl)



trimethoxysilane (MPTS) and then, the thiol groups were oxidized to sulfonic acid by hydrogen peroxide.

The TGA analysis of SBA-Pr-SO<sub>3</sub>H (Fig. 4) confirmed the amount of organic groups on SBA-15. The weight reduction of SBA-Pr-SO<sub>3</sub>H in the temperature range between 200-600°C indicated that the amount of organic group was 1.2 mmol/g.

The nitrogen adsorption-desorption isotherms for SBA-15 and SBA-Pr-SO<sub>3</sub>H are shown in Fig. 5. Both materials show type-IV adsorption behavior with the hysteresis loops appearing at relatively high pressure, suggesting that the prepared samples have regular mesoporous framework structures. Their texture properties are given in Table 2. The surface area, average pore diameter calculated by the BET method and pore volume of SBA-Pr-SO<sub>3</sub>H are 440 m<sup>2</sup>g<sup>-1</sup>, 6.0 nm and 0.660 cm<sup>3</sup> g<sup>-1</sup>, respectively, which are smaller than those

**Table 2. Porosimetry values for SBA-15 and functionalized SBA-15.**

	Surface area (cm <sup>2</sup> g <sup>-1</sup> )	Pore volume (cm <sup>3</sup> g <sup>-1</sup> )	Pore diameter (nm)
SBA-15	649	0.806	6.2
SBA-SO <sub>3</sub> H	440	0.660	6.0

**Fig. 4: Thermogravimetric analysis (TGA) of SBA-Pr-SO<sub>3</sub>H.****Fig. 5: N<sub>2</sub> adsorption-desorption isotherms and pore size distribution (inset) for SBA-15 (---) and SBA-Pr-SO<sub>3</sub>H (—).**

of SBA-15 due to the immobilization of sulfonosilane groups into the pores.

## CONCLUSIONS

In conclusion, an efficient synthesis of polyhydroquinolines derivatives has been achieved via a one-pot four component reaction of aromatic aldehydes, dimedone,  $\beta$ -ketoesters, and ammonium acetate under

solvent free conditions using the reusable and environmentally benign sulfonic acid functionalized nanoporous SBA-15 as solid acid catalyst. The reasonable reaction time, very good to excellent yields, simple workup procedure, and environmentally friendly conditions are main merits of this method. Therefore SBA-Pr-SO<sub>3</sub>H can be used as an efficient heterogeneous catalyst such as other catalysts in organic syntheses [33].

## Acknowledgements

We gratefully acknowledge for financial support from the Research Council of Alzahra University and University of Tehran.

Received : Oct. 2, 2008 ; Accepted : July 29, 2009

## REFERENCES

- [1] (a) Godfraid T., Miller R., Wibo M., Calcium Antagonism and Calcium Entry Blockade, *Pharmacol. Rev.* **38**, p. 321 (1986);  
(b) Sausins A., Duburs G., Synthesis of 1,4-Dihydropyridines by Cyclocondensation Reactions, *Heterocycles*, **27**, p. 269 (1988);  
(c) Mager P.P., Coburn R.A., Solo A.J., Triggler D.J., Rothe H., QSAR, Diagnostic Statistics and Molecular Modelling of 1,4-Dihydropyridine Calcium Antagonists: A Difficult Road Ahead, *Drug Des. Discovery*, **8**, p. 273 (1992);  
(d) Mannhold R., Jablonka B., Voigt W., Schonafinger K., Schraven K., Calcium- and Calmodulin-Antagonism of Elnadipine Derivatives: Comparative SAR, *Eur. J. Med. Chem.* **27**, p. 229 (1992).
- [2] (a) Klusa V., Cerebrocrast, Neuroprotectant, Cognition Enhancer, *Drugs Future*, **20**, p. 135 (1995);  
(b) Bretzel R.G., Bollen C.C., Maester E., Federlin K.F., Nephroprotective Effects of Nitrendipine in Hypertensive Type I and Type II Diabetic Patients, *Am. J. Kidney Dis.*, **21**, p. 54 (1993);  
(c) Boer R., Gekeler V., Chemosensitizers in Tumor Therapy: New Compounds Promise Better Efficacy, *Drugs Future*, **20**, p. 499 (1995).
- [3] (a) Hantzsch A., Mittheilungen Synthese von Thiazolen und Oxazolonen, *Ber. Dtsch. Chem. Ges.* **21**, p. 942, (1888);  
(b) Wiley R.H., England D.C., Behr L.C., "In Organic Reactions", John Wiley, **6**, p. 367 (1951);

- (c) Dondoni A., Massi A., Minghini E., Bertolasi V., Multicomponent Hantzsch Cyclocondensation as a Route to Highly Functionalized 2- and 4-Dihydropyridylalanines, 2- and 4-Pyridylalanines, and Their N-Oxides: Preparation Via a Polymer-Assisted Solution-Phase Approach, *Tetrahedron* **60**, p. 2311 (2004).
- [4] (a) Khadikar B.M., Gaikar V.G., Chitnavis A.A., Aqueous Hydrotrope Solution as a Safer Medium for Microwave Enhanced Hantzsch Dihydropyridine Ester Synthesis, *Tetrahedron Lett.* **36**, p. 8083 (1995); (b) Ohberg L., Westman J., An Efficient and Fast Procedure for the Hantzsch Dihydropyridine Synthesis Under Microwave Conditions, *Synlett* 1296 (2001); (c) Agarwal A., Chauhan P.M.S., Solid Supported Synthesis of Structurally Diverse Dihydropyrido [2,3-d] Pyrimidines Using Microwave Irradiation, *Tetrahedron Lett.*, **46**, p. 1345 (2005).
- [5] (a) Ji S.-J., Jiang Z.-Q., Lu J., Loh T.-P., Facile Ionic Liquids-Promoted One-Pot Synthesis of Polyhydroquinoline Derivatives Under Solvent Free Conditions, *Synlett*, **831** (2004); (b) Sridhar R., Perumal P.T., A New Protocol to Synthesize 1,4-Dihydropyridines by Using 3,4,5-Trifluorobenzeneboronic Acid as a Catalyst in Ionic Liquid: Synthesis of Novel 4-(3-Carboxyl-1H-Pyrazol-4-yl)-1,4-Dihydropyridines, *Tetrahedron*, **61**, p. 2465 (2005).
- [6] (a) Phillips A.P., Hantzsch's Pyridine Synthesis, *J. Am. Chem. Soc.* **71**, p. 4003 (1949); (b) Anderson G.J.R., Berkelhammer G., A Study of the Primary Acid Reaction on Model Compounds of Reduced Diphosphopyridine Nucleotide, *J. Am. Chem. Soc.*, **80**, p. 992 (1958); (c) Singh H., Chimni D.S.S., Kumar S., Acid Catalysed Enamine Induced Transformation of 1,3-Dimethyl-5-Formyluracil. A Unique Annulation Reaction with Enaminones, *Tetrahedron*, **51**, p. 12775 (1995); (d) Gordeev M.F., Patel D.V., Gordon E.M., Approaches to Combinatorial Synthesis of Heterocycles: A Solid-Phase Synthesis of 1,4-Dihydropyridines, *J. Org. Chem.* **61**, p. 924 (1996); (e) Breitenbucher J.G, Figliozzi G., Solid-Phase Synthesis of 4-aryl-1,4-Dihydropyridines via the Hantzsch Three Component Condensation, *Tetrahedron Lett.*, **41**, p. 4311, (2000); (f) Liang J.-C., Yeh J.-L., Wang C.-S., Liou S.-F., Tasi C.-H., Chen I.-J., The New Generation Dihydropyridine Type Calcium Blockers, Bearing 4-Phenyl Oxypropanolamine, Display  $\alpha$ - $\beta$ -Adrenoceptor Antagonist and Long-Acting Antihypertensive Activities, *Bioorg. Med. Chem.*, **10**, p. 719, (2002); (g) Miri R., Niknahad H., Vesal Gh., Shafiee A., Synthesis and Calcium Channel Antagonist Activities of 3-Nitrooxyalkyl, 5-Alkyl 1,4-Dihydro-2,6-Dimethyl -4-(1-Methyl-5-Nitro-2-Imidazolyl)-3, 5-Pyridinedicarboxylates, *IL Farmaco*, **57**, p. 123 (2002); (h) Dondoni A., Massi A., Minghini E., Sabbatini S., Bertoasi V., Model Studies Toward the Synthesis of Dihydropyrimidinyl and Pyridyl  $\alpha$ -Amino Acids Via Three-Component Biginelli and Hantzsch Cyclocondensations, *J. Org. Chem.*, **68**, p. 6172, (2003); (i) Dondoni A., Massi A., Minghini E., Bertoasi V., Multicomponent Hantzsch Cyclocondensation as a Route to Highly Functionalized 2- and 4-Dihydropyridylalanines, 2- and 4-Pyridylalanines, and Their N-Oxides: Preparation Via a Polymer-Assisted Solution-Phase Approach, *Tetrahedron*, **60**, p. 2311 (2004); (j) Tewari N., Dwivedi N., Tripathi R.P., Tetrabutylammonium Hydrogen Sulfate Catalyzed Eco-Friendly and Efficient Synthesis of Glycosyl 1,4-Dihydropyridines, *Tetrahedron Lett.*, **45**, p. 9011 (2004); (k) Moseley J.D., Alternative Esters in the Synthesis of ZD0947, *Tetrahedron Lett.*, **46**, p. 3179 (2005).
- [7] Sabitha G., Reddy G.S.K.K., Reddy Ch.S., Yadav J.S., A Novel TMSI-Mediated Synthesis of Hantzsch 1,4-Dihydropyridines at Ambient Temperature, *Tetrahedron Lett.*, **44**, 4129 (2003).
- [8] Wang L.-M., Sheng J., Zhang L., Han J.-W., Fan Z., Tian H., Qian C.-T., Facile Yb(OTf)<sub>3</sub> Promoted One-Pot Synthesis of Polyhydroquinoline Derivatives Through Hantzsch Reaction, *Tetrahedron*, **61**, p. 1539 (2005).
- [9] Ko S., Sastry M.N.V., Lin C., Yao C., Molecular Iodine-Catalyzed One-Pot Synthesis of 4-Substituted-1,4-Dihydropyridine Derivatives Via Hantzsch Reaction, *Tetrahedron Lett.*, **46**, p. 5771 (2005).



- [10] Beck S., Vartuli J.C., Roth W.J., Kresge C.T., Leonowicz M.E., Schmitt K.D., Chu C.T.-W., Olson D.H., Sheppard E.W., McCullen S.B., Higgins J.B., Schlenker J.L., A New Family of Mesoporous Molecular Sieves Prepared with Liquid Crystal Templates, *J. Am. Chem. Soc.* **114**, p. 10834, (1992).
- [11] Reinert P., Garcia B., Morin C., Badiei A., Perriat P., Tillement O., Bonneviot L., Cationic Templating with Organic Counterion for Superstable Mesoporous Silica, Nanotechnology in Mesoporous Materials, *Stud. Surf. Sci. Catal.*, **146**, p. 133, (2003).
- [12] Bonneviot L., Morin M., Badiei A., Mesoporous Metal or Nano-Metal Oxides and Method for Making Same, Patent No. 01/55031 A1, (2001).
- [13] Zhao D., Huo Q., Feng J., Chmelka B.F., Stucky G.D., Nonionic Triblock and Star Diblock Copolymer and Oligomeric Sulfonate Syntheses of Highly Ordered, Hydrothermally Stable, Mesoporous Silica Structures, *J. Am. Chem. Soc.*, **120**, p. 6024 (1998).
- [14] Trong On D., Desplandier-Giscard D., Danumah C., Kaliaguine S., Perspectives in Catalytic Applications of Mesoporous Materials, *Appl. Catal. A: Gen.*, **222**, p. 299 (2001).
- [15] Mohammadi Ziarani G., Badiei A., Miralami A., The Study of Solvent Effect on the Diastereoselectivity of Diels-Alder Reaction in the Presence of Nanoporous Silica-Supported Cerium Sulfonate Catalyst, *Eur. J. Sci. Res.*, **18**, p. 282 (2007).
- [16] Ganjali M.R., Daftari A., Hajiagha Babaei L., Badiei A., Saberyan K., Mohammadi Ziarani G., Moghimi A., Pico Level Monitoring of Silver with Modified Hexagonal Mesoporous Compound (MCM-41) and Inductively Coupled Plasma Atomic Emission Spectrometry, *Water, Air, Soil Pollut.*, **173**, p. 71, (2006).
- [17] Ganjali M.R., Hajiagha Babaei L., Badiei A., Mohammadi Ziarani G., Tarlani A., Novel Method for the Fast Preconcentration and Monitoring of a ppt Level of Lead and Copper with a Modified Hexagonal Mesoporous Silica Compound and Inductively Coupled Plasma Atomic Emission Spectrometry, *Anal. Sci.*, **20**, p. 725, (2004).
- [18] Ganjali M.R., Hajiagha Babaei L., Badiei A., Saberian K., Behbahani S., Mohammadi Ziarani G., Salavati-Niasari M., A Novel Method for Fast Enrichment and Monitoring of Hexavalent and Trivalent Chromium at the PPT Level with Modified Silica MCM-41 and its Determination by Inductively Coupled Plasma Optical Emission Spectrometry, *Quim. Nova*, **29**, p. 440 (2006).
- [19] Badiei A., Norouzi P., Tousi F., Study of Electrochemical Behavior and Adsorption Mechanism of  $[\text{Co}(\text{en})_2\text{Cl}_2]^+$  on Mesoporous Modified Carbon Paste Electrode, *Eur. J. Sci. Res.*, **12**, p. 39 (2005).
- [20] Zhang H.-X., Cao A.-M., Hu J.-S., Wan L.-J., Lee S.-T., Electrochemical Sensor for Detecting Ultratrace Nitroaromatic Compounds Using Mesoporous  $\text{SiO}_2$ -Modified Electrode, *Anal. Chem.*, **78**, p. 1967 (2006).
- [21] Walcarius A., Despas C., Bessiere J., Selective Monitoring of Cu(II) Species Using a Silica Modified Carbon Paste Electrode, *Anal. Chim. Acta*, **385**, p. 79, (1999).
- [22] Lim M.H., Blanford C.F., Stein A., Synthesis of Ordered Microporous Silicates with Organosulfur Surface Groups and Their Applications as Solid Acid Catalysts, *Chem. Mater.* **10**, p. 467 (1998).
- [23] Wight A.P., Davis M.E., Design and Preparation of Organic-Inorganic Hybrid Catalysts, *Chem. Rev.*, **102**, p. 3589 (2002).
- [24] a) van Rhijn W.M., De Vos D., Sels B.F., Bossaert W.D., Jacobs P.A., Sulfonic Acid Functionalised Ordered Mesoporous Materials as Catalysts for Condensation and Esterification Reactions, *Chem. Commun.*, **317** (1998);  
b) Das B., Venkateswarlu K., Holla H., Krishnaiah M., Sulfonic Acid Functionalized Silica: A Remarkably Efficient Heterogeneous Reusable Catalyst for  $\alpha$ -Monobromination of Carbonyl Compounds Using *N*-Bromosuccinimide, *J. Mol. Catal. A: Chem.* **253**, p. 107 (2006);  
c) Onaka M., Hashimoto N., Kitabata Y., Yamasaki R., Aluminum-Rich Mesoporous Aluminosilicate (Al-HMS) as a Solid Acid Catalyst for the Diels-Alder Reaction of Acrylates with 1,3-Dienes, *Appl. Catal. A: General.*, **241**, p. 307 (2003).
- [25] Kureshy, R. I., Ahmad, I., Pathak, K., Khan, N. H., Abdi, S. H. R., Jasra, R. V., Sulfonic Acid Functionalized Mesoporous SBA-15 as an Efficient and Recyclable Catalyst for the Synthesis of Chromenes from Chromanols, *Catal. Commun.*, **10**, p. 572 (2009).

- [26] Karimi B., Zareyee D., Design of a Highly Efficient and Water-Tolerant Sulfonic Acid Nanoreactor Based on Tunable Ordered Porous Silica for the von Pechmann Reaction, *Org. Lett.*, **10**, p. 3989 (2008).
- [27] Nagarapu L., Kumari M.D., Kumari N.V., Kantevari S., MCM-41 Catalyzed Rapid and Efficient One-Pot Synthesis of Polyhydroquinolines Via the Hantzsch Reaction Under Solvent-Free Conditions, *Catal. Commun.*, **8**, p. 1871 (2007).
- [28] Mohammadi Ziarani G., Badiei A., Miralami A., A Study of the Diastereoselectivity of Diels-Alder Reactions on the Ce-SiO<sub>2</sub> as Support, *Bull. Korean Chem. Soc.* **29**, p. 47 (2008).
- [29] a) Kumar A., Maurya R.A., Synthesis of Polyhydroquinoline Derivatives Through Unsymmetric Hantzsch Reaction Using Organocatalysts, *Tetrahedron*, **63**, p. 1946 (2007);  
b) Kumar A., Maurya R.A., Bakers Yeast Catalyzed Synthesis of Polyhydroquinoline Derivatives via an Unsymmetrical Hantzsch Reaction, *Tetrahedron Lett.* **48**, p. 3887 (2007).
- [30] Heravi M., Bakhtiari Kh., Javadi N.M., Bamoharam F., Saeedi M., Oskooie H., K<sub>7</sub>[PW<sub>11</sub>CoO<sub>40</sub>]-Catalyzed One-Pot Synthesis of Polyhydroquinoline Derivatives via the Hantzsch Three Component Condensation, *J. Mol. Catal. A*, **264**, p. 50 (2007).
- [31] Donelson L.J., Gibbs R.A., De S., An Efficient One-Pot Synthesis of Polyhydroquinoline Derivatives Through the Hantzsch Four Component Condensation, *J. Mol. Catal. A*, **256**, p. 309 (2006).
- [32] Song G., Wang B., Wu X., Kang Y., Yang L., Montmorillonite K10 Clay: An Effective Solid Catalyst for One-Pot Synthesis of Polyhydroquinoline Derivatives, *Synth. Commun.*, **35**, p. 2875 (2005).
- [33] a) Shaabani A., Maleki A., A Fast and Efficient Method for the Synthesis of 1,5-Benzodiazepine Derivatives Under Solvent-Free Conditions, *Iran J. Chem. Chem. Eng.*, **26**, p. 93 (2007);  
b) Bigdeli M.A., Nahid N., Heravi M.M., Sulphuric Acid Adsorbed on Silica Gel. A Remarkable Acetylation Catalyst, *Iran J. Chem. Chem. Eng.*, **19** (2000).